

7-1-2008

Psychopathy and Adolescent Females: Does Gender Alter the Relation Between Childhood Trauma and PCL:YV Scores?

Trevor Barese

Roger Williams University, tbarese@hotmail.com

Follow this and additional works at: http://docs.rwu.edu/psych_thesis



Part of the [Psychology Commons](#)

Recommended Citation

Barese, Trevor, "Psychopathy and Adolescent Females: Does Gender Alter the Relation Between Childhood Trauma and PCL:YV Scores?" (2008). *Psychology Theses*. Paper 17.

http://docs.rwu.edu/psych_thesis/17

This Thesis is brought to you for free and open access by the Feinstein College of Arts and Sciences Theses at DOCS@RWU. It has been accepted for inclusion in Psychology Theses by an authorized administrator of DOCS@RWU. For more information, please contact mwu@rwu.edu.

Psychopathy and Adolescent Females: Does Gender Alter the Relation Between
Childhood Trauma and PCL:YV Scores?

Thesis submitted in partial fulfillment of the requirements for the degree
Master of Arts in Forensic Psychology

Feinstein College of Arts and Sciences
Roger Williams University

Trevor Barese

Class of 2008

July 24, 2008

Program Director's Signature

Thesis Advisor's Signature

Abstract

A growing rate of violence among adolescent females has led to an increased interest in gender differences associated with the assessment and development of psychopathy. This study utilized a sample of 100 youthful offenders, to examine the role of gender in the relation between various forms of childhood trauma and scores on the Psychopathy Checklist: Youth Version (PCL:YV). Correlations were found between certain forms of childhood abuse and elevated PCL:YV scores. Furthermore, gender appeared to alter the predictive power of proposed developmental risk factors commonly associated with psychopathy. The findings indicate possible discrepancies in developmental risk factors between genders. Implications for guiding intervention and treatment strategies are discussed.

Acknowledgements

I would like to take this opportunity to acknowledge the individuals who made this project possible. First, I would like to express my gratitude to Dr. Frank DiCataldo who served as my committee chair. It was Dr. DiCataldo who initially got me thinking about possible research topics. This project would not have been possible without his assistance in accessing Bedford Policy Institute's database and gaining approval from the Department of Youth Services to use the case file information. I would also like to acknowledge Dr. Donald Whitworth. Dr. Whitworth has had a tremendous impact on my academic development. His passion for psychology and appreciation for the scientific method have been a continuing source of inspiration. Dr. Whitworth provided thoughtful feedback and much needed emotional support through the thesis process. In addition, I would like to express my gratitude to Dr. Gina Vincent for agreeing to serve as an external member of my committee. Dr. Vincent was extremely helpful in providing an outside perspective on the study. Her extensive knowledge in the area of psychopathy was an invaluable source of feedback. Finally, I would like to acknowledge my thesis partner, Nathan Cook. Nate's attention to detail, particularly deadlines, was crucial in moving this project forward. His level of dedication and high academic standards are beyond reproach. More important, his shared passion for the development of novel and meaningful research made this an exciting and enjoyable process.

Dedication

I would like to dedicate this thesis to my family and loved ones. Particularly my mother, Dr. Evelyn Barese who has been a continuing source of inspiration throughout my life. Her integrity and unwavering dedication to the pursuit of knowledge and intellectual-growth continue to amaze me. Most important, she has taught me never to compromise my beliefs and values but also to maintain an open mind. Without her unconditional love and support I would have never made it this far.

I would also like to thank the Psychology department at Roger Williams University, especially Dr. Whitworth. Dr. Whitworth has taught me a great deal about the human psyche and the importance of being genuine. I truly admire him as an educator and clinician.

Psychopathy and Adolescent Females: Does Gender Alter the Relation Between Childhood Trauma and PCL:YV Scores?

The psychopath, according to Cleckley, is characterized by a lack of: responsibility, honesty, sincerity, guilt or shame, capacity for deep attachment, and insight into his condition (1976). In addition, the psychopath is egotistical and appears unable to learn from his past transgressions. Although, psychopathy is not included in the Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision (DSM-IV TR) it is nevertheless recognized as a legitimate personality disorder within the field of psychology (Hemphill & Hart, 2003). First identified by Philip Pinel in 1801, as an emotional pathology, this conceptual disorder has since been studied by social and medical scientists under a variety of names for over two centuries. Yet, it was not until 1915 that Emil Kraepelin coined the term “psychopathic personality”, in reference to a group of criminals seeming to lack a sense of morals (Lykken, 1996). Cleckley later defined psychopathic individuals as, “hotheaded; cold-hearted; impulsive; irresponsible; selfish; emotionally shallow; manipulative; and lacking in empathy, anxiety, and remorse” (Lynam, 1999). In *The Mask of Sanity*, Cleckley eloquently illustrates the psychopath’s lack of emotional intelligence, stating “Beauty and ugliness, except in a very superficial sense, goodness, evil, love, horror, and humor have no actual meaning, no power to move him” (1976, p. 40).

Perhaps the psychopathic personality has captured the interest of so many scientists due to the relative rarity of the disorder and the severity of its impact on society. Despite higher rates within the prison population (15-25%), base rates for psychopathy are generally very low (Hare, 2003). In fact, less than one percent of the

general adult population would meet the diagnostic criteria for psychopathy (Hare). Though small, this group of individuals is believed to be responsible for a disproportionately large amount of crime and resulting monetary loss to society (Hare, McPherson, & Forth, 1988; Hart, Kropp, & Hare, 1988; Moffitt, 1993). Psychopathic individuals are at an increased risk of reoffending and doing so: more quickly, more often, and more violently than non-psychopathic offenders (Salekin, Rogers, & Sewell, 1996). In a comprehensive study by David Anderson (1999) the annual burden of crime in the United States was estimated in excess of one trillion dollars. If, as the literature suggests, psychopathic individuals do in fact commit more crimes, of a greater variety and over a longer span of time (Forth & Burke, 1998) then it can be inferred that they create a disproportionately large financial strain on tax payers and society in general.

Accordingly, the ability to identify adolescents who might be at increased risk of developing psychopathic personalities would be invaluable. In general adolescents are more likely to engage in criminal behaviors than are adults. Rather than being pathological, in certain settings “antisocial” behaviors might be considered normative among adolescent groups (Vincent & Grisso, 2005). Therefore, it is not “adolescent-limited” offenders, those who exhibit late onset antisocial behavior over a brief period of time, but rather the “life-course-persistent” offenders, those who have exhibited antisocial behaviors from very early on and seem unable to desist, that are most closely linked to psychopathy and of the greatest concern to society (Moffitt, 1993).

Identification of specific childhood risk factors associated with the development of psychopathic or life-course-persistent antisocial adult personalities are crucial preliminary steps in the advancement of effective screening and intervention strategies

(Forth & Mailloux, 2000). Detection of empirically supported risk factors will assist in the recognition of youths at elevated risk of developing personalities conducive to criminal and antisocial lifestyles. The accurate identification of these youths will facilitate the more efficient and effective implementation of intervention strategies (Forth & Mailloux; Frick, 2001; 2004). Thus, the continued study of at-risk youth and possible etiological factors associated with psychopathy are ethically and financially viable pursuits. Although the importance of such research has not gone unnoticed within the field of psychology it has, until fairly recently, focused primarily on male adolescent offenders.

Current Trends in Female Offending

Over the last couple of decades, the United States Department of Justice has reported a significant increase in the arrest and imprisonment rates of adult and adolescent females. According to a survey by Snell and Morton (1991), female arrest rates increased by 24% and the number of females incarcerated increased by 75% between 1986 and 1991. Yet, during this five year span male arrest rates only increased by 13% and male imprisonment rates increased by 53%. Interestingly, the 1998 per capita arrest rate among juvenile females was nearly twice that of adult females (Greenfeld & Snell, 2000). As of 2006 the number of female prisoners under state or federal jurisdiction was 112,498, a 33% increase from 1998 (Greenfeld & Snell; Sabol, Couture, & Harrison, 2007). What's more, the percentage of the prison population accounted for by females has risen from 4.7% in 1986 to 7% in 2006 (Sabol et al.; Snell & Morton). Despite the growing rate of female involvement within the legal system, the majority of psychopathy literature has relied heavily on samples of male offenders

(Odgers & Moretti, 2002; Warren et al., 2003). This lack of literature, specific to female offending, has left social scientists to rely heavily on findings obtained from male samples in guiding clinical judgments and treatment strategies.

Assessment of Psychopathy among Adolescents

The growing rate of violence among females and subsequent involvement in the legal system combined with the recent downward application of the psychopathy construct has led to an increased interest in the valid assessment of psychopathy among adolescent females (Odgers & Moretti, 2002; Odgers, Reppucci, & Moretti, 2005). Historically the assessment of psychopathy in adolescents and children has relied primarily on modified forms of adult assessment tools (Forth et al., 2003). Unfortunately, these tools relied mostly on Factor 2 (Socially Deviant Lifestyle) traits, or overt antisocial behaviors (Forth et al.). More recently, formal assessment tools have been developed with the primary goal of assessing psychopathy among children and adolescents (Frick & Hare, 2001; Hare & Hervé, 1999; Lilienfeld & Widows, 2005; Lynam, 1997). In response to growing research demands and the inadequacies of available assessment tools, modified versions of the Psychopathy Checklist Revised (PCL-R; Hare, 1991; 2003) began to be utilized by researchers. This led to the development and publication of the Psychopathy Checklist: Youth Version (PCL: YV; Forth, Kosson, & Hare, 2003), as a research tool for assessing psychopathy during adolescents. The majority of research pertaining to adolescent females has relied largely on the PCL-R, and the PCL:YV. The validity of these tools in the prediction of future offending and violence has been supported through a growing number of studies

(Corrado, Vincent, Hart, & Cohen, 2004; Forth & Burke, 1998; Gretton, Hare, & Catchpole, 2004; Guy, Edens, Anthony, & Douglas, 2005).

Collectively, the literature supports at least three general hypotheses, of particular relevance to the current study: adult psychopathic females differ significantly from males in prevalence, factor structure, and comorbid diagnosis; childhood trauma in the forms of abuse and unstable environment have been positively correlated with total PCL scores, and at least two subtypes of psychopathy exist. These hypotheses and their implications will be discussed in further detail later.

Precautions

Since its introduction the PCL:YV has been used extensively to research adolescent psychopathy in relation to youth violence and recidivism. The motivating force behind these studies is the hope of early intervention and prevention of adult psychopathy. However, the stigma attached to the label of psychopathy has caused much trepidation within the field of psychology as well as the juvenile courts (Forth et al., 2003). More important, there is still a great deal of research needed to address the relation between gender and PCL:YV scores (Odgers, Reppucci et al., 2005).

It is important to note that under no circumstance should psychopathy be considered a disorder diagnosable during childhood or adolescence. Rather, psychopathy is a personality disorder, and by definition must reflect a stable pattern of behavior that is present over a significant portion of an individual's life (American Psychiatric Association, 2000; Zaitchik & Barese, 2008). The identification and examination of adolescents that share behaviors similar to those of adult psychopaths should be aimed at gaining insight into possible developmental processes and in no way as an attempt to

diagnose or label adolescents as psychopathic. Regardless of ethical and theoretical concerns, no tool currently exists capable of the reliable and therefore valid assessment of future psychopathy among adolescents.

The little longitudinal research that has been done does not provide overwhelming support for the global stability of psychopathic traits from adolescents into adulthood (Frick, Kimonis, Dandreaux, & Farell, 2003). For instance, a longitudinal study comparing assessments of psychopathy at age 13 with follow up assessments of the same group at age 24 found only moderate correlations between childhood and adult total scores ($r^2 = .31$) (Lynam, Caspi, Moffitt, Loeber, & Stouthamer-Loeber, 2007). This study illustrates the limitations of attempting to assess psychopathy during childhood and early adolescents. To label a child psychopathic at age thirteen would be unwise and unethical, given that roughly 70% of children appearing psychopathic at age 13 would not meet the criteria for diagnosis, a decade later, at age 24. The tools currently available for the assessment of psychopathy in adolescents continue to lack the predictive validity necessary to outweigh the risk of a false-positive diagnosis. Thus, it would be unethical and unfounded to extend the assessment of psychopathy among adolescents beyond the realm of empirical research (see Frick, 2002; Hart, Watt, & Vincent, 2002; Lynam, 2002; Seagrave & Grisso, 2002; Skeem & Cauffman, 2003).

Gender Differences

The literature supports gender differences in: factor structure of the PCL-R and PCL:YV, prevalence of psychopathy and associated concurrent comorbid diagnosis for both adult and adolescent samples (Forouzan & Cooke, 2005; Nicholls, Ogloff, Brink, & Spidel, 2005; Strand & Belfrage, 2005). These differences suggest that the construct of

psychopathy, largely developed using male samples, may differ between genders in: etiology, expression, and prognosis (Odgers, Moretti, & Reppucci, 2005). If females who score high on the PCL display unique factor loadings this might suggest membership in a distinct subgroup of psychopathy or the presence of an unidentified, confounding personality disorder.

Differences in the prevalence of psychopathy between genders have been found in a large number of studies using different versions of the Psychopathy Checklist (Grann, 2000; Vitale & Newman, 2001). According to Nicholls and Petrila, base rates of adult psychopathy in female offender samples range between 7.5 and 23% compared to 15 and 30% in male offender samples (2005). However, precise differences in base rate are difficult to assess due to a lack of standardized cut off scores for female samples between studies. A number of researchers have used lower cut off scores, such as 25, when assessing psychopathy among females in an attempt to compensate for lower prevalence rates and mean total scores (Forouzan & Cooke, 2005; Vitale & Newman). Unfortunately, these procedural modifications contribute to the difficulty in interpretation of findings and the ambiguity surrounding differences in prevalence between genders.

Generally, female subjects obtain lower total scores on the PCL-R in comparison to males (Forth, Brown, Hart, & Hare, 1996; Grann, 2000; Vitale, Smith, Brinkley, & Newman, 2002). For instance, using the suggested cut off score of 18 on the PCL:SV, Strand and Belfrage (2005) found total prevalence rates of 16% among adult female offenders and 25% among adult male offenders. Similarly, in a study of 103 adult female offenders only 16% scored above the cutoff score of 29 on the PCL-R (Salekin, Rogers,

& Sewell, 1997). This lower prevalence rate among females has likely contributed to an overrepresentation of males in the psychopathy literature (Schrum & Salekin, 2006).

In addition to differences in prevalence between genders, research implies that females exhibit PCL-R factor structures distinct from those of males (Grann, 2000; Jackson, Rogers, Neumann, & Lambert, 2002). Forouzan and Cooke state, “A core requirement of gender equivalence is that the factor structures should be equivalent” (2005, pp.769). Yet research, using the 2 factor model, has found males to consistently load higher on Factor 2 (Socially Deviant Lifestyle) of the PCL-R than do females with similar total scores (Schrum & Salekin, 2006). Furthermore, some of the literature has failed to support the validity of the two factor model among female samples. As a result the majority of recent psychopathy studies utilizing female samples have relied on the three or four factor models (Strand & Belfrage, 2005). These findings bring in to question the construct validity of psychopathy among females.

Similar to the discrepancies in factor structure a few studies have examined the convergent validity of psychopathy between genders in the prevalence of concurrent comorbid psychological disorders among psychopaths (Odgers et al. 2007; Vitale et al., 2002; Warren et al., 2003). These studies examine gender differences between psychopaths on types and rates of personality disorder. If psychopathy is essentially the same between genders, one would expect to find similar constellations of comorbid diagnoses between genders. Differences in comorbid personality disorders are to be expected when taking in to account factor structure, and item response differences between genders (Strand & Belfrage, 2005).

Childhood Risk Factors

The literature has identified a number of social or environmental factors positively correlated with PCL-R, PCL:YV scores, and psychopathy in general. As early as the 1950s and 60s sociologists and psychologists have expressed an interest in the parent-child relationships of sociopathic or psychopathic individuals. Early on, Hare (1970) described how studies utilizing semantic differential procedures implied that psychopathic or sociopathic adults evaluated their parents more negatively than other subjects. He explained how inconsistent or postponed discipline might facilitate the development of psychopathic personalities in individuals with higher physiological thresholds of anticipatory anxiety. Further, he theorized that separation from primary caregivers was related to the development of psychopathy. However, at that time “psychopathy” as a personality disorder had not been clearly operationalized and as a result lacked construct validity and therefore reliability of assessment. Consequently, many studies relied on self-report measures such as the Maudsley Personality Inventory (MPI) or the Psychopathic Deviate (Pd) scale from the Minnesota Multiphasic Personality Inventory (MMPI; Hare, 1970).

Almost three decades later, in a review of the literature surrounding early childhood trauma and family background as they relate to PCL:YV scores, Forth and Burke (1998) provide support for findings supportive of Hare’s theory. Poor discipline, poor school experience, and parental rejection were found to be significant predictors of PCL-R total scores, with inconsistent parenting found to be the strongest predictor of Factor 1 (Interpersonal\Affective) scores. Notably, lack of parental supervision and parental rejection were the two variables most strongly linked to psychopathy throughout all of the studies (Forth & Burke). In an unpublished study by Burke and Forth (1996),

the authors created a global scale of family background variables in order to assess the combined influence on PCL:YV scores. Their findings indicate that negative family experiences are associated with increased total, Factor 1 (interpersonal + affective), and Factor 2 (behavioral + antisocial) scores on the PCL:YV. However, these findings were not significant among young offenders (in Forth & Burke; see also Forth & Mailloux, 2000).

More recently, Lynam and Gudonis (2005) offered an overview of theoretical models of development designed to explain these correlations. One such theory implicates childhood abuse and disrupted early attachments as key risk factors associated with the development of psychopathy (Saltaris, 2002). For example, foster care placement, a form of disrupted attachment, and history of physical abuse have been positively correlated with PCL:YV scores (Campbell, Porter, & Santor, 2004). Additionally, both childhood physical and sexual abuse have been linked to high total PCL-R scores (Marshall & Cooke, 1995; Verona, Hicks, & Patrick, 2005; Weiler & Widom, 1996). Although statistically significant, these risk factors are generally weak predictors of psychopathy, typically accounting for only a small amount of variance in total scores (Poythress, Skeem, & Lilienfeld, 2006). When interpreting these findings it is important to consider the likely covariance between childhood abuse and disruptions of attachment and living situations.

The literature linking childhood trauma to criminal behavior and lifestyles is seemingly corroborated by the crime statistics offered by the Bureau of Justice. For example, in 1991, 42% of female and 43% of male inmates reported having been raised by a single parent. In addition 17% of both male and female inmates reported ever living

in a foster home, agency, or institution while growing up (Snell & Morton, 1991). Despite similarities between genders in reported disruptions of family structure, gender differences do appear to exist in reported levels of physical and sexual abuse. The percentages of female offenders and male offenders who reported ever being physically or sexually abused were 43% and 12%, respectively. Thirty two percent of females compared to 11% of males reported being abused prior to age 18. Females reported relatively equal instances of physical and sexual abuse, about 32% whereas males reported twice the amount of physical abuse (10%) as sexual abuse (5%). Interestingly, female offenders who were the victims of abuse were more likely than non-abused female offenders to be in prison for a violent offense (42% v. 25%) and less likely to be serving a sentence for either a drug offense (25% v. 38%) or a property offense (25% v. 31%). Of all violent female offenders those who had experienced abuse were significantly more likely to be sentenced for homicide (Snell & Morton). This information is relevant to the study, in that psychopathy has been empirically linked to increased rates of violent crime.

In general the findings suggest the existence of multiple developmental pathways to becoming a psychopathic adult (Forth & Burke, 1998). These pathways are likely guided by a combination of both environmental and biological factors. It is important to note, all of these studies have relied on retrospective reports of childhood trauma. Thus, caution must be taken when making inferences as to causation.

Psychopathy Subtypes

In 1948 Karpman separated psychopathy into two subgroups, “primary psychopathy” and “secondary psychopathy”. Primary psychopathy involved personality

traits such as, “callousness, manipulativeness, glibness, and lack of anxiety and remorse” (Lynam, 1999). Secondary psychopathy dealt with the stress-related antisocial patterns of behavior of psychopaths (Lynam). It was not until 1991 that Hare operationalized these two subcategories, renaming them “Factor 1” and “Factor 2”. Factor 1 was defined as “a personality style associated with (the) callous; remorseless exploitation of others”; Whereas Factor 2 involved “an impulsive, unstable, antisocial lifestyle” (Kosson & Kelly, 1997). This new distinction maintained the underlying differences originally proposed by Karpman.

The existence of psychopathic subtypes have been revealed using a number of personality tests to assess the concurrent validity of the PCL-R and PCL:YV and identify within group heterogeneity using cluster analysis (Falkenbach, Poythress, & Creevy, 2008; Murphy & Vess, 2003; Vassileva, Kosson, Abramowitz, & Conrod, 2005; Vincent, Vitacco, Grisso, & Corrado, 2003). Subtypes have been developed not only to explain apparent behavioral differences between psychopaths but also as a bases for proposing unique etiological models and differentiating risk of recidivism (Poythress et al., 2006). The two subtypes of psychopathy known as: primary or emotionally stable psychopathy and secondary or aggressive psychopathy, have been supported using model-based cluster analysis of PCL-R and Multidimensional Personality Questionnaire in brief form (MPQ-BF) scores. Primary psychopathy has been characterized by low stress reaction, and increased level of control or planning; whereas, secondary psychopathy has been characterized by aggressive behavior and high stress reaction (Hicks, Markon, Newman, Patrick, & Krueger, 2004). The overly emotional and impulsive secondary psychopath doesn't fit Hervey Cleckley's (1976) classic conceptualization of psychopathy.

Conversely, the research suggests that secondary psychopaths comprise the majority of adult males diagnosed with psychopathy (Hicks et al.).

Despite a paucity of research on subtype prevalence among adolescent females, findings pertaining to factor structure indicate that female factor loading on the PCL-R is more similar to that of males within the primary psychopathy group (Hicks et al., 2004; Jackson et al., 2002). Thus, a possible discrepancy in subtype prevalence between genders might account for gender base rate differences among adult psychopaths. Furthermore, subtypes may present distinct etiologies and thereby mediate the developmental pathway and degree to which early environmental factors are predictive of future PCL-R and PCL:YV scores. The reliable and valid assessment of gender-specific risk factors associated with the development of psychopathy is crucial to creating effective early intervention and prevention strategies for at-risk youth.

Hypotheses:

1. The following forms of childhood abuse will be correlated with high PCL:YV scores: physical abuse, sexual abuse, emotional abuse, and neglect.
2. PCL:YV scores will be positively correlated with number of previous living arrangements.
3. PCL:YV scores will be negatively correlated with age at first interruption of family structure.
4. Positive parental support or nurturance and parental control or accountability will be negatively correlated with PCL:YV total scores

5. PCL:YV factor loading, for the two factor model, will differ between genders. Females will obtain higher Factor 1 scores and lower Factor 2 scores in comparison to males.
6. Early childhood abuse, unstable living arrangement, and early interruption of family structure will predict PCL:YV scores.
7. Gender will alter the degree to which early childhood abuse, unstable living arrangement, and early interruption of family structure are predictive of PCL:YV scores.

Method

Subjects

This study utilized a sample of 100 youthful offenders, consisting of 50 females and 50 males, ranging in age from 13 to 19 years old ($M = 16$ years; $SD = 1.2$). All subjects were committed to a state juvenile justice agency in the northeastern United States. The mean age of subjects did not differ significantly between genders (females $M = 15.8$; $SD = 1.2$; males $M = 16.3$; $SD = 1.2$). It is important to note the mode for males was 17 years old, comprising 52% of the entire male sample. Approximately one half of the sample was Caucasian (49%) and one quarter African American (26%). The remaining subjects consisted of 18% Hispanic, 4% biracial, and 3% Asian. There were no significant differences in the distribution of race between gender groups (Table 1).

All of the subjects were evaluated by the Forensic Evaluation Service of the Bedford Policy Institute between the years of 1996 and 2003. Upon a request from the Massachusetts Department of Youth Services, the Bedford Policy Institute developed, implemented, and operated an evaluation service designed to assess risk and treatment needs of juvenile offenders. The Forensic Evaluation Service began in 1996 and by

2003, had completed approximately 2,800 evaluations compiling an extensive computer database of juvenile records. All evaluations were conducted by doctoral-level psychologists, licensed in Massachusetts, and possessing the added credential of Designated Forensic Psychologist (DFP) by the Massachusetts Department of Mental Health. Subjects were randomly selected from the database by a Bedford Policy Institute employee, blind to the hypotheses of the current study. All forensic evaluation reports included in the sample had the subjects' names and other identifiers redacted and replaced with an identifying number. Thus, the identities of participating offenders were kept strictly confidential. Data about each youthful offender was collected solely from case files and forensic mental health reports. There were no attempts to contact subjects. The study strictly followed American Psychological Association ethical guidelines as well as relevant policies set forth by the Massachusetts Department of Youth Services Institutional Review Board and the Roger Williams University Human Subject Review Board.

Materials

The Psychopathy Checklist: Youth Version (PCL:YV).

The youth version of the PCL maintains essentially the same 20 items of the PCL-R (Appendix A). However, the items have been modified to fit the experiences and social expectations unique to adolescents. For instance items pertaining to: marital relationships, occupational history, and past criminality were rephrased and rescaled accordingly. Each item is rated on a three point scale (0 = No, 1 = Maybe, 2 = Yes) there is also an option to omit an item. The sum of all 20 items provides a total score ranging

between 0 and 40. If items were omitted prorated scores are available (Forth et al., 2003).

Unlike the PCL-R, no cut off scores have been provided for use with the PCL: YV (Forth et al., 2003). However, the research indicates fairly parallel distributions of total scores between the youth and adult versions, with adolescents' generally scoring about five points lower than adults. In response, much of the research with adolescents has used total scores of 25 and above to represent high-Psychopathy groups (Forth et al.). Much like the PCL-R, two, three, and four factor models are available for use with the PCL: YV to examine both, individual and group loading differences (Jones, Cauffman, Miller, & Mulvey, 2006). The current study utilized both the two and four factor models to explore possible discrepancies in factor loading between genders and distinct variable correlations. Both the two and four factor models consist of the same 18 items (see Appendix B). The four factor model includes four item clusters: F1: Interpersonal (4 items), F2: Affective (4 items), F3: Behavioral (5 items), F4: Antisocial (5 items). The two factor model combines both, F1 and F2 to form a single factor, Factor 1 (Interpersonal/Affective), as well as F3 and F4 to form a single factor, Factor 2 (Socially Deviant Lifestyle). The two factor model is thought to capture the key features associated with primary (Factor 1) and secondary (Factor 2) psychopathy subtypes.

Archival information from subjects' case files was used to obtain scores on the Psychopathy Checklist: Youth Version (PCL:YV; see Appendix C). Valid scoring the PCL:YV solely through the use of archival data has been empirically supported (Guy & Douglas, 2006) and permitted in the technical manual (Forth, 2005; Forth et al., 2003). The PCL:YV is made up of twenty items that are scored as either 0 if the trait does not

apply to the youth, 1 if the trait is present but not to a substantial degree, or 2 if the trait is definitely present; the maximum score on the PCL:YV is 40. To aid in the scoring and determination of each trait the evaluator is provided with an item description and some behavioral examples. The inter-rater reliability of the PCL:YV has been supported, with a single-rater intra-class correlation ranging from .90 to .96 (Forth et al.).

Each subject, in the study, was scored by one of two trained raters. Thirty (30%N) cases were randomly selected to be scored independently by both raters in order to assess and establish adequate inter-rater reliability of PCL:YV total and factor scores. To establish inter-rater reliability, Intra-class Correlation Coefficients (ICC) were computed for PCL:YV: total scores (ICC = .95), Factor 1: Interpersonal/Affective scores (ICC=.97), Factor 2: Socially Deviant Lifestyle scores (ICC= .85), F1: Interpersonal scores (ICC = .91), F2: Affective scores (ICC = .94), F3: Behavioral scores (ICC = .83), and F4: Antisocial scores (ICC = .86). These results maintain an acceptable level of agreement between raters implying that subjects were scored reliably between raters.

Procedure

The case information used in this study was part of a computer database compiled through the Forensic Evaluation Service of the Bedford Policy Institute. The evaluations completed as part of this service were comprehensive and extensive. Initial assessments included: a full review of relevant records and reports, consultations with caseworkers, team members, and program clinicians, as well as a complete and thorough clinical interview. The clinical interviews focused on historical risk factors and the youth's current level of functioning and goals for the future. During the interview juveniles were asked to provide an account of their past, highlighting shifts in family structure,

memorable experiences, and social support networks. On the basis of the material gathered, an evaluation was prepared with the intent of informing and aiding the classification of offenders and the identification of relevant treatment needs. To avoid experimenter bias, subjects' family histories were omitted during scoring of the PCL:YV. Permission to utilize the case information was gained through the Massachusetts Department of Youth Services and Bedford Policy Institute (see Appendix D).

Upon completion of the forensic evaluation, the evaluator, or other trained Bedford Policy Institute employee, extracted information relevant to six broad areas and coded it on a forensic evaluation data sheet (FEDS; see Appendix E). The six areas represented on the data sheet include: 1) demographic information (e.g., age, gender, etc.); 2) delinquency history information (i.e., list of prior delinquency adjudication and legal findings); 3) mental health history and data (e.g., prior psychiatric hospitalization, current medication, history of suicide attempts, etc.); 4) clinical data/risk factors (e.g., history of abuse, substance abuse problems, mode of violence); 5) nature of the offense (e.g., age of victim, gender of victim, relationship to victim, etc.); and 6) clinical judgments (e.g., type of service recommended, risk factors identified, treatment needs, etc.). The information from the data sheet was then entered into a computer database. With the exception of page 2 (delinquency adjudication and legal findings) this information was omitted during scoring of the PCL:YV and used in conjunction with family histories to code the independent variables.

Variables

For this study, the variables considering subjects' histories of abuse, attachment problems, parental support or control, and exposure to domestic violence were coded

with the aid of the FEDS (see Appendix E). All variables were coded by the same licensed psychologist who authored the subject's evaluation for DYS. These were recorded by the researcher from the FEDS. Permission to utilize the forensic evaluation data sheets was gained through the Massachusetts Department of Youth Services and Bedford Policy Institute. There were seven items of primary interest. These are described below.

Positive Parental Support or Nurturance

“Positive Parental Support or Nurturance” was a Clinical Data / Risk Factor. The evaluator was provided three response choices: yes, no, or not clear. In order to reduce possible ambiguities only yes or no responses were used in the final analysis. All variables originally coded as not clear were recoded as missing data.

Parental Control and Accountability for Juvenile

“Parental Control and Accountability for Juvenile” was recorded as: yes, no, or not clear (see Appendix E). In order to reduce possible ambiguities only yes or no responses were used in the final analysis. All variables originally coded as not clear were recoded as missing data.

History of Attachment Problems Early Childhood

“History of attachment problems early childhood” was recorded by the evaluators as provided three response choices: yes, no, or not clear (see Appendix E). In order to reduce possible ambiguities only yes or no responses were used in the final analysis. All variables originally coded as not clear were recoded as missing data.

History of Abuse

Upon completion of each assessment the evaluator completed the FED form, which includes “History of abuse” under section IV (Clinical Data / Risk Factors). The evaluator was provided two response choices: yes or no (see Appendix E). In order to establish the reliability of this variable all answers were checked against the subject’s files to corroborate the finding.

Type of Abuse

Type of abuse was categorized as: physical, sexual, emotional, and neglect. Evaluators were allowed to provide multiple responses to this item, representing multiple forms of abuse (see Appendix E). In order to establish the reliability of this variable all answers were checked against the subject’s files to corroborate these findings.

Prior History of DSS Services

Upon completion of each assessment the evaluator completed the FED form, which includes “Prior History of DSS Services” under section IV (Clinical Data / Risk Factors). The evaluator was provided two response choices: yes or no (see Appendix E). This item was coded solely on the evaluator’s response to this question.

Witnessed Domestic Violence

Upon completion of each assessment the evaluator completed the FED form, which includes “Witnessed domestic violence” among a list of factors, under section VI (Conclusions / Risk factors identified). The evaluator simply checked all items that were applicable (see Appendix E). Therefore, all subjects who received a check next to “Witnessed domestic violence” were coded as yes and all others were coded as no. In order to establish the reliability of this variable all answers were checked against the subject’s files to corroborate these findings.

Number of Previous Living Arrangements

In addition to the previously coded variables found in subjects' FED forms, each subject was also coded on two factors at the time of the study. Subjects' "Number of Previous Living Arrangements" was coded, using case files, by summing the number of shifts in family structure our guardian (see Appendix F). For example if a subject lived at home with his biological family was placed in foster care for a year and returned home to his biological family he would be coded as a three. All shifts in living arrangement were coded up until the juveniles governing offense. Transitions between secure facilities following the juvenile's governing offense were not included. Additionally parental separation and acquisition of live-in significant others, step-parents, or step-siblings were defined as shifts in living arrangement. This item was coded separately from PCL:YV, after all PCL:YVs had been scored. Additionally, the family background section of each case file was not used in the scoring of the PCL:YV.

Age at First Interruption of Family Structure

Subjects' "Age at First Interruption of Family Structure" was coded, using case files, according to the earliest age (year) at which they experienced a significant shift in family structure (see Appendix F). For example if a subject lived at home with her biological mother from birth until age three when she was sent to live with her grandmother, she would be coded as a three. If the subject did not experience any early interruption of family structure prior to their commitment to DYS, then the age at which they entered DYS was coded. This item was coded separately from PCL:YV, after all PCL:YVs had been scored. Additionally, the family background section of each case file was not used in the scoring of the PCL:YV.

Results

Descriptive Statistics

PCL:YV scores did not differ significantly between genders (see Table 1). Additionally, PCL:YV total scores did not differ significantly between races, though African American subjects did obtain higher scores than Caucasian subjects, means of 19.4 and 16.6 respectively. Total PCL:YV scores for female subjects ranged from 6 to 30, with a mean of 18 (SD = 5.2). Total PCL:YV scores for male subjects ranged from 8 to 35, with a mean of 17.7 (SD = 6.1; Table 1). There were no significant differences between genders on the two factor model for either Factor 1 (Interpersonal/Affective) scores (M = 7.1, SD = 3.5), or Factor 2 (Socially Deviant Lifestyle) scores (M = 9.1, SD = 2.7). Similarly, factor scores did not differ significantly between genders on the four factor model. Using the four factor model, subjects obtained the following average factor scores; F1: Interpersonal (M = 3.0, SD = 2.1); F2: Affective (M = 4.1, SD = 2.1); F3: Behavioral (M = 4.5, SD = 1.6); F4: Antisocial (M = 4.5, SD = 1.8).

The majority of the sample had experienced some form of abuse (71%), 84% of female subjects and 58% of male subjects had a history of abuse. A majority of the sample (67%) had contact with the Department of Social Services (DSS) prior to their commitment offense. The mean age at first interruption of family structure was 8.9 years (SD = 5.3). No significant differences in age at first interruption of family structure were found between genders (see Table 1). The mean number of previous living arrangements for the entire sample was 4.4 (SD = 4.3). The average number of previous living arrangements did not differ significantly between genders (see Table 1).

A chi square analysis revealed significant differences in history of abuse between genders, $\chi^2 (1, N = 100) = 8.2, p = .004$, with females being more likely to have experienced abuse. In fact, females were twice as likely as males to have been either sexually abused $\chi^2 (1, N = 100) = 6.3, p = .012$ or emotionally abused $\chi^2 (1, N = 100) = 5.5, p = .019$. Females were significantly more likely than males to have had prior involvement with the Department of Social Services (DSS) $\chi^2 (1, N = 100) = 5.5, p = .019$.

Hypothesis 1: The following forms of childhood abuse will be correlated with high PCL:YV scores: physical abuse, sexual abuse, emotional abuse, and neglect.

The hypothesized correlation between childhood abuse and PCL:YV scores was partially supported. Chi square analysis revealed significant relations between high PCL:YV scores and history of sexual abuse $\chi^2 (1, n = 62) = 5, p = .025$ (see Table 2). Subjects with total PCL:YV scores above 21 had significantly higher rates of sexual abuse compared to those with total scores below 15. Similarly the relation between high total PCL:YV scores and neglect approached significance $\chi^2 (1, n = 62) = 3.4, p = .065$ (see Table 2). Subjects who obtained high scores on the PCL:YV had higher rates of childhood neglect. Chi squares were performed for history of abuse, history of physical abuse and history of emotional abuse and high and low PCL:YV groups. The results failed to support the hypothesis, that physical and emotional abuse would be significantly correlated with PCL:YV scores. Interestingly, prior involvement with DSS was significantly correlated with high PCL:YV scores $\chi^2 (1, n = 62) = 4.3, p = .039$ (see Table 2).

Hypothesis 2: PCL:YV scores will be positively correlated with number of previous living arrangements.

Pearson product-moment correlations failed to support the hypothesis that, PCL:YV total scores would be positively correlated with number of previous living arrangements. Number of previous living arrangements was not significantly correlated with PCL:YV total scores. In addition, number of previous living arrangements was not significantly correlated with either Factor 1 (Interpersonal/Affective) or Factor 2 (Socially Deviant Lifestyle) scores. Furthermore, splitting the sample by gender did not result in a change of significance.

Hypothesis 3: PCL:YV scores will be negatively correlated with age at first interruption of family structure.

Pearson product-moment correlations failed to support the hypothesis that, PCL:YV total scores would be negatively correlated with age at first interruption of family structure. Age at first interruption of family structure was not significantly correlated with PCL:YV total scores. In addition, age at first interruption of family structure was not significantly correlated with either Factor 1 (Interpersonal/Affective) or Factor 2 (Socially Deviant Lifestyle) scores. Furthermore, splitting the sample by gender did not result in a change of significance.

Hypothesis 4: Positive parental support or nurturance and parental control or accountability will be negatively correlated with PCL:YV total scores

Chi square analysis supported the hypothesis that, parental control and accountability would be negatively correlated with high PCL:YV total scores, $\chi^2(1, n = 53) = 5.6, p = .017$. Subjects who scored above 21 on the PCL:YV were less likely to

have had parental control or accountability. The presence of positive parental support or nurturance was not significantly correlated with high PCL:YV scores. Next, the sample was split by gender and the chi square analyses were repeated for both positive parental support or nurturance and parental control and accountability with PCL:YV total scores. Females scoring below 15 on the PCL:YV were more likely to have had parents who provided positive support and nurturance than females who obtained high (>21) scores, $\chi^2(1, n = 22) = 4.8, p = .029$. Whereas, males scoring below 15 on the PCL:YV were more likely to have had parents who provided adequate control and accountability than males obtaining high (>21) scores $\chi^2(1, n = 27) = 4.2, p = .04$ (see Table 3).

Hypothesis 5: PCL:YV factor loading, for the two factor model, will differ between genders. Females will obtain higher Factor 1 scores and lower Factor 2 scores in comparison to males.

Multiple regressions partially supported the hypothesis that, early childhood abuse, unstable living arrangements, and early interruption of family structure would act as predictors of PCL:YV scores. In order to uncover the best predictors of PCL:YV total scores, standard multiple regressions were performed. The best-fitting model, accounting for 12% of the total variance, in PCL:YV scores, revealing: “Number of Previous Living Arrangements”, “Prior History of DSS Services”, and “Parental Control and Accountability for Juvenile” as significant predictors of subjects’ PCL:YV total scores: $F(3, 83) = 3.9, p = .012$. The best predictor of PCL:YV total score within this model was “Parental Control and Accountability for Juvenile” ($\beta = -.25, p = .021$). Subjects who were viewed as having parental control and accountability were more likely to obtain low scores on the PCL:YV, whereas subjects who were perceived as lacking

parental control and accountability were more likely to obtain high scores on the PCL:YV. Prior History of DSS Services was a significant predictor of PCL:YV total score ($\beta = .23, p = .048$). Finally, “Number of Previous Living Arrangements” did not contribute significantly to the model as an independent variable ($\beta = -.12, p = .29$) (see Table 4). Additional regressions suggest that histories of: abuse, physical abuse, emotional abuse, witnessing domestic violence, and attachment problems were not significant predictors of PCL:YV total score for the sample as a whole.

Hypothesis 6 Early childhood abuse, unstable living arrangement, and early interruption of family structure will predict PCL:YV scores.

Using the two factor model, Independent samples t-tests failed to support significant differences between genders on mean Factor 1 (Interpersonal/Affective) and mean Factor 2 (Socially Deviant Lifestyle) scores. Both males and females scored approximately 2 points higher on Factor 2 (mean = 9) in comparison to Factor 1 (mean = 7). To further investigate the possibility of differences in subtype prevalence rates between genders a categorical variable was created based on differences of factor scores within subjects. All subjects obtaining a Factor 1 (Interpersonal/Affective) score greater than their Factor 2 (Socially Deviant Lifestyle) score were coded as primary and all subjects obtaining a Factor 2 score greater than their Factor 1 score were coded as secondary. All subjects displaying equal scores on both Factor 1 and Factor 2 were excluded. After splitting the sample by gender and excluding all subjects with PCL:YV total scores < 20 a simple analysis of frequency for subtype was performed. Despite the small sample size, subtype did not appear to differ between genders. However, about two thirds of both males and females who scored above 20 on the PCL:YV fell into the

secondary group ($n_f = 11$, $n_m = 10$). Only one third of both males and females who scored above 20 on the PCL:YV fell into the primary group ($n_f = 5$, $n_m = 6$).

Hypothesis 7: Gender will alter the degree to which early childhood abuse, unstable living arrangement, and early interruption of family structure are predictive of PCL:YV scores.

Multiple regressions partially support the influence of gender on the degree to which early childhood abuse, unstable living arrangement, and early interruption of family structure are predictive of PCL:YV scores. To uncover the degree to which gender might alter the relation between childhood trauma and PCL:YV total scores the sample was split by gender and the above regression (Number of Previous Living Arrangements, Prior History of DSS Services, and “Parental Control and Accountability for Juvenile as predictors of PCL:YV total score) was repeated. After splitting the sample by gender the model was no longer significant for female subjects. However, the model increased in effect size when used exclusively for predicting PCL:YV total scores of male subjects. This model, accounting for 19% of the total variance, using “Number of Previous Living Arrangements”, “Prior History of DSS Services”, and “Parental Control and Accountability for Juvenile” as significant predictors of male subjects’ PCL:YV total scores, $F(3, 38) = 3.0$, $p = .044$. The only significant independent predictor of PCL:YV total score within this model was Prior History of DSS Services ($\beta = .38$, $p = .043$). Finally, “Parental Control and Accountability for Juvenile” ($\beta = -.23$, $p = .13$) and “Number of Previous Living Arrangements” did not contribute significantly to the model on their own ($\beta = -.18$, $p = .31$) (see Table 5).

Further multiple regressions imply inconsistencies between genders in terms of which variables are significant predictors of PCL:YV total and factor scores. For example, “number of previous living arrangements”, “history of sexual abuse”, and “positive parental support or nurturance” were significant predictors of female Factor 1 (Interpersonal/Affective) scores. However these variables were not significant predictors of PCL:YV Factor 1 (Interpersonal/Affective) scores among male subjects.

In order to uncover the best predictors of PCL:YV Factor 1 (Interpersonal / Affective) scores, standard multiple regressions were performed. The best-fitting model, accounting for 25.8% of the total variance in PCL:YV Factor 1 (Interpersonal/Affective), revealed “number of previous living arrangements”, “history of sexual abuse”, and “positive parental support or nurturance” as significant predictors of female subjects’ PCL:YV Factor 1 (Interpersonal/Affective) scores, $F(3, 34) = 3.9, p = .016$. The single best predictor of PCL:YV Factor 1 (Interpersonal/Affective) score in this model was “positive parental support or nurturance”, which explained 19% of the total variance ($\beta = -.46, p = .006$). Thus, females who were viewed as having positive parental support or nurturance were more likely to obtain low scores on Factor 1 (Interpersonal/Affective) of the PCL:YV and females who were perceived as lacking positive parental support or nurturance were more likely to obtain high scores on Factor 1 (Interpersonal/Affective) of the PCL:YV. Similarly, “number of previous living arrangements” was predictive of female PCL:YV Factor 1 (Interpersonal/Affective) score, accounting for 9% of the total variance ($\beta = -.31, p = .056$). Finally, “history of sexual abuse” did not contribute significantly to the model as an independent variable ($\beta = .16, p = .31$) (see Table 6).

Discussion

This study examined the role of gender in the relation between childhood trauma and PCL:YV scores. In general the results support correlations between certain childhood traumas and increased PCL:YV scores. Specifically, childhood sexual abuse and neglect were correlated with high PCL:YV scores. Whereas, childhood physical and emotional abuse were not associated with high PCL:YV scores. Multiple regressions supported the number of previous living arrangements, a history of DSS involvement, and the presence of parental control or accountability, as predictors of PCL:YV total score. It is important to note, history of DSS may have been a strong predictor because it captured a group of subjects who likely experienced significant and confirmed instances of sexual abuse and neglect. Both of which were significantly correlated with high PCL:YV total scores. Essentially, subjects who had experienced sexual abuse or neglect would be more likely to have had involvement with DSS in response to these experiences. Although these variables were significant predictors, as suggested by the literature they accounted for a rather small amount of the variance in PCL:YV total score.

As hypothesized, gender did appear to alter the relation between proposed risk factors and PCL:YV scores. The presence of positive parental support or nurturance was significantly correlated with low PCL:YV scores for females, but not males. However, the presence of parental control or accountability was correlated with low PCL:YV scores for males, but not females. The importance of positive parental support and nurturance for females, and parental control and accountability for males, as possible protective factors suggests that effective intervention strategies may differ between genders.

Furthermore, multiple regressions failed to support the universality of predictor variables between genders on both total and two factor PCL:YV scores. The presence of

gender-specific predictors suggests that there may be discrepancies in developmental risk factors between genders. Interestingly, the number of previous living arrangements, a history of sexual abuse, and the presence of positive parental support or nurturance were significant predictors of females' Factor 1 (Interpersonal/Affective) scores. Though insignificant, females did score higher than males on Factor 1 and lower than males on Factor 2. In addition, female subjects were twice as likely to have experienced sexual or emotional abuse compared to male subjects. Thus, it would seem plausible that a history of sexual or emotional abuse might lead to the development of Post Traumatic Stress Disorder (PTSD). As a result, symptoms of PTSD may mimic affective and interpersonal traits commonly associated with psychopathy, leading to inflated Factor 1 scores on the PCL:YV.

Limitations

This study was limited in that all data was gathered archivally, which limited the amount of control over what information was available for scoring and coding. Additionally, the sample was rather small and very few subjects obtained total scores above 25 on the PCL:YV, especially after splitting the sample by gender. The mean PCL:YV total score for the entire sample was 18. This is relatively low in comparison to the typical mean total score of 24 obtained in other studies of institutionalized adolescents. In addition, there was a high rate of abuse among the sample. In fact, the majority of subjects (71%) had experienced some form of abuse.

Moreover, subtype differences between genders were difficult to assess in this study and interestingly as total scores increased the number of primary females significantly declined. All four females who scored above 25 on the PCL:YV were

coded as secondary. Two of the four males who scored above 25 on the PCL:YV were coded as primary and the other two were coded as secondary. Therefore it would be unwise to make any assumptions about subtype prevalence based on a small sample of relatively low scoring individuals. Furthermore the method of classification used in designating subtypes was overly simplistic and may not have accurately captured the key characteristics associated with each of the subtype groups.

One limitation of this study is possible within group heterogeneity in terms of age at onset of antisocial behavior. The PCL:YV alone does not provide an indication of the age at which subjects began their antisocial behavior. According to Moffitt and Caspi, there is a distinct difference between antisocial adolescents with early onset in comparison to those with onset beginning at adolescents, in relation to etiological factors (2001). Children with early onset tend to exhibit a more stable and nonmalleable pattern of antisocial behavior similar to that of the adult psychopath. These are the individuals who seem most effected by early childhood trauma and parental attachment issues. Furthermore, Moffitt and Caspi suggest a greater difference in prevalence rates of life-course persistent offenders between genders in comparison to adolescent limited offenders. Their research suggests a life-course persistent ratio of 10:1 and an adolescent-limited ratio of 1.5:1 for males and females respectively (Moffitt & Caspi). The failure to discriminate between these two groups may confound the relation between early childhood experiences and the development of adult psychopathic personalities.

Implications

Differences in modes of expressed violence between genders have been empirically supported among adults and adolescents who have obtained high scores on

the PCL (Odgers, Moretti et al., 2005). These differences are often viewed as different outlets for the same disorder rather than distinct outlets for distinct disorders. If females who exhibit psychopathic traits differ significantly from males not only in etiology and distinct developmental pathways, but also in the manner of manifestation and mode of expression, perhaps they should be viewed as having a distinct disorder. Any attempt at effective identification, intervention, and treatment aimed at youth considered high risk for the development of antisocial or psychopathic personality disorders must be framed around the development and symptoms of the disorder. The findings of this study and the existing literature suggest that distinctions be made between genders in the identification of risk factors and resulting treatment needs.

Finally, the failure of longitudinal studies in the early identification of adult psychopathy should not be viewed as a failure or a shortcoming but as a sign that personalities are malleable during adolescence. Rather than discouragement these findings should provide motivation in the development of effective intervention strategies. It has been debated whether antisocial and criminal behaviors are a necessary product of psychopathy or an unpleasant but avoidable side effect. Yet, many individuals who appear to share the affective and interpersonal characteristics associated with psychopathy manage to adapt their behaviors to societal expectations by finding socially acceptable lifestyles congruent to their unique character traits. Through the continued study of adolescents at risk of developing psychopathic personalities and antisocial lifestyles it is hoped that we can effectively intervene in this process; Thereby increasing the quality of life for both the adolescent and society as a whole.

*Hyp 6 These results did not support differences in subtype prevalence between genders. However, due to the small sample size, relatively low cutoff score, and rudimentary procedure of subtype classification, few inferences should be made from these findings.

References

- American Psychiatric Association. (2000). *Diagnostic and Statistical Manual of Mental Disorders*, Fourth Edition, Text Revision. Washington, DC: APA.
- Anderson, D. A. (1999). The aggregate burden of crime. *Journal of Law and Economics*, 42 (2), 611-642.
- Campbell, M. A., Porter, S., & Santor, D. (2004). Psychopathic traits in adolescent offenders: An evaluation of criminal history, clinical, and psychosocial correlates. *Behavioral Sciences & the Law*, 22, 23-47.
- Cleckley, H. (1976). *The Mask of Sanity* (5th ed.). St. Louis, MO: Mosby.
- Corrado, R., Vincent, G., Hart, S., & Cohen, I. (2004). Predictive validity of the Psychopathy Checklist: Youth Version for general and violent recidivism. *Behavioral Sciences & the Law*, 22(1), 5-22.
- Falkenbach, D., Poythress, N., & Creevy, C. (2008). The exploration of subclinical psychopathic subtypes and the relationship with types of aggression. *Personality and Individual Differences*, 44 (4), 821-832.
- Forouzan, E., & Cooke, D. (2005). Figuring out la femme fatale: Conceptual and assessment issues concerning psychopathy in females. *Behavioral Sciences & the Law*, 23(6), 765-778.
- Forth, A. (2005). Hare Psychopathy Checklist: Youth Version. *Mental health screening and assessment in juvenile justice* (pp. 324-338). New York, NY, US: Guilford Press.
- Forth, A. E., Brown, S. L., Hart, S. D., & Hare, R. D. (1996). The assessment of

psychopathy in male and female noncriminals: Reliability and validity.

Personality and Individual Differences, 20, 531-543.

Forth, A.E., & Burke, H. (1998). Psychopathy in adolescence; Assessment, violence, and developmental precursors. *Psychopathy; Theory, Research, and Implications for Society*, 205-229.

Forth, A.E., Kosson, D.S., & Hare, R.D. (2003). *The Hare Psychopathy Checklist: Youth Version*. Toronto: Multi-Health Systems.

Forth, A., & Mailloux, D. (2000). Psychopathy in youth: What do we know? *The Clinical and Forensic Assessment of Psychopathy: A Practitioner's Guide* (pp. 25-54). Mahwah, NJ, US: Lawrence Erlbaum Associates Publishers.

Frick, P. J. (2001). Effective interventions for children and adolescents with conduct disorder. *The Canadian Journal of Psychiatry*, 46, 597-608

Frick, P. (2002). Juvenile psychopathy from a developmental perspective: Implications for construct development and use in forensic assessments. *Law and Human Behavior*, 26(2), 247-253.

Frick, P. J. (2004). Developmental pathways to conduct disorder: Implications for serving youth who show severe aggressive and antisocial behavior. *Psychology in the Schools*, 41 (8), 823-834.

Frick, P. J., & Hare, R. D. (2001). *The Antisocial Process Screening Device (APSD)*. Toronto, ON: Multi-Health Systems.

Frick, P., Kimonis, E., Dandreaux, D., & Farrell, J. (2003). The 4 year stability of psychopathic traits in non-referred youth. *Behavioral Sciences & the Law*, 21(6), 713-736.

- Grann, M. (2000). The PCL-R and gender. *European Journal of Psychological Assessment, 16*(3), 147-149.
- Greenfeld, L.A., & Snell, T.L. (2000). Women offenders. *Bureau of Justice Statistics Bulletin*.
- Gretton, H., Hare, R., & Catchpole, R. (2004). Psychopathy and offending from adolescence to adulthood: A 10-year follow-up. *Journal of Consulting and Clinical Psychology, 72*(4), 636-645.
- Guy, L., & Douglas, K. (2006). Examining the utility of the PCL:SV as a screening measure using competing factor models of psychopathy. *Psychological Assessment, 18* (2), 225-230.
- Guy, L., Edens, J., Anthony, C., & Douglas, K. (2005). Does psychopathy predict institutional misconduct among adults? A meta-analytic investigation. *Journal of Consulting and Clinical Psychology, 73*(6), 1056-1064.
- Hare, R.D. (1970). *Psychopathy Theory and Research*. U.S.A.: John Wiley & Sons, Inc.
- Hare, R.D. (1991). *The Hare Psychopathy Checklist – Revised*. Toronto: Multi-Health Systems.
- Hare, R.D. (2003). *The Hare Psychopathy Checklist – Revised: 2nd Ed*. Toronto: Multi-Health Systems.
- Hare, R.D., & Hervé, H.F. (1999). Hare P-Scan: Research Version Manual. Multi-Health Systems: Toronto.
- Hare, R., McPherson, L., & Forth, A. (1988). Male psychopaths and their criminal careers. *Journal of Consulting and Clinical Psychology, 56*(5), 710-714.
- Hart, S., Kropp, P., & Hare, R. (1988). Performance of male psychopaths following

- conditional release from prison. *Journal of Consulting and Clinical Psychology*, 56(2), 227-232.
- Hart, S., Watt, K., & Vincent, G. (2002). Commentary on Seagrave and Grisso: Impressions of the state of the art. *Law and Human Behavior*, 26(2), 241-245.
- Hemphill, J., & Hart, S. (2003). Forensic and clinical issues in the assessment of psychopathy. *Handbook of Psychology: Forensic Psychology, Vol. 11* (pp. 87-107). Hoboken, NJ, US: John Wiley & Sons Inc.
- Hicks, B. M., Markon, K. E., Newman, J. P., Patrick, C. J., & Krueger, R. F. (2004). Identifying psychopathy subtypes on the basis of personality structure. *Psychological Assessment*, 16 (3), 276-288.
- Jackson, R. L., Rogers, R., Neumann, C. S., & Lambert, P. L. (2002). Psychopathy in female offenders: An investigation of its underlying dimensions. *Criminal Justice and Behavior*, 29, 692-704.
- Jones, S., Cauffman, E., Miller, J. D., & Mulvey, E. (2006). Investigating different factor structures of the psychopathy checklist: Youth version: Confirmatory factor analytic findings. *Psychological Assessment*, 18, 33-48.
- Kosson, D.S., & Kelly, J.C. (1997). Psychopathy-related traits predict self-reported sexual aggression among college men. *Journal of Interpersonal Violence*, 12, 241-255.
- Lilienfeld, S. O., & Widows, M. (2005). *Psychopathic Personality Inventory-Revised professional manual*. Odessa, FL: PAR.
- Lorber, M. F. (2004). Psychophysiology of aggression, Psychopathy, and conduct problems: a meta-analysis. *Psychological Bulletin*, 130, 531-552.

- Lykken, D.T. (1996). Psychopathy, sociopathy, and crime. *Society*, 34, 29-39.
- Lynam D.R. (1997). Pursuing the psychopath: Capturing the fledgling psychopath in a nomological net. *Journal of Abnormal Psychology* 106: 425-438.
- Lynam, D.R. (1999). Self-reported Psychopathy: A validation study. *Journal of Personality Assessment*, 73, 110-123.
- Lynam, D. (2002). Fledgling psychopathy: A view from personality theory. *Law and Human Behavior*, 26(2), 255-259.
- Lynam, D., Caspi, A., Moffitt, T., Loeber, R., & Stouthamer-Loeber, M. (2007). Longitudinal evidence that Psychopathy scores in early adolescence predict adult Psychopathy. *Journal of Abnormal Psychology*, 116(1), 155-165.
- Lynam, D. R., & Gudonis, L. (2005). The development of psychopathy. *Annual Review of Clinical Psychology*, 1, 381-407.
- Lynam, D., Loeber, R., & Stouthamer-Loeber, M. (2008). The stability of psychopathy from adolescence into adulthood: The search for moderators. *Criminal Justice and Behavior*, 35(2), 228-243.
- Marshall, L., & Cooke, D. (1995). The role of childhood experiences in the etiology of psychopathy. *Issues in Criminological & Legal Psychology*, 24, 107-108.
- Moffitt, T. E. (1993). Adolescence-limited and life-course-persistent antisocial behavior: A developmental taxonomy. *Psychological Review*, 100 (4), 674-701.
- Moffitt, T., & Caspi, A. (2001). Childhood predictors differentiate life-course persistent and adolescence-limited antisocial pathways among males and females. *Development and Psychopathology*, 13(2), 355-375.
- Murphy, C., & Vess, J. (2003). Subtypes of psychopathy: Proposed differences between

- narcissistic, borderline, sadistic, and antisocial psychopaths. *Psychiatric Quarterly*, 74(1), 11-29.
- Nicholls, T., Ogloff, J., Brink, J., & Spidel, A. (2005). Psychopathy in women: A review of its clinical usefulness for assessing risk for aggression and criminality. *Behavioral Sciences & the Law*, 23(6), 779-802.
- Nicholls, T. L., Petrila, J. (2005). Gender and psychopathy: An overview of important issues and introduction to the special issue. *Behavioral Sciences and the Law*, 23, 729-741.
- Odgers, C., & Moretti, M. (2002). Aggressive and antisocial girls: Research update and challenges. *International Journal of Forensic Mental Health*, 1(2), 103-119.
- Odgers, C., Moretti, M., Burnette, M., Chauhan, P., Waite, D., & Reppucci, N. (2007). A latent variable modeling approach to identifying subtypes of serious and violent female juvenile offenders. *Aggressive Behavior*, 33(4), 339-352.
- Odgers, C., Moretti, M., & Reppucci, N. (2005). Examining the science and practice of violence risk assessment with female adolescents. *Law and Human Behavior*, 29(1), 7-27.
- Odgers, C. L., Reppucci, N. D., & Moretti, M. M. (2005). Nipping psychopathy in the bud: An examination of the convergent, predictive, and theoretical utility of the PCL-YV among adolescent girls. *Behavioral Sciences & the Law*, 23, 743-763.
- Poythress, N., Skeem, J., & Lilienfeld, S. (2006). Associations among early abuse, dissociation, and psychopathy in an offender sample. *Journal of Abnormal Psychology*, 115(2), 288-297.
- Sabol, W. J., Couture, H., & Harrison, P.M. (2007). Prisoners in 2006. *Bureau of Justice*

Statistics Bulletin.

- Salekin, R., Rogers, R., & Sewell, K. (1996). A review and meta-analysis of the Psychopathy Checklist and Psychopathy Checklist-Revised: Predictive validity of dangerousness. *Clinical Psychology: Science and Practice*, 3(3), 203-215.
- Salekin, R., Rogers, R., & Sewell, K. (1997). Construct validity of psychopathy in a female offender sample: A multitrait-multimethod evaluation. *Journal of Abnormal Psychology*, 106(4), 576-585.
- Saltaris, C. (2002). Psychopathy in juvenile offenders: Can temperament and attachment be considered as robust developmental precursors? *Clinical Psychology Review*, 22, 729-752.
- Schrum, C. L., & Salekin, R. T. (2006). Psychopathy in adolescent female offenders: An item response theory analysis of the psychopathy checklist: Youth version. *Behavioral Sciences & the Law*, 24, 39-63.
- Seagrave, D., & Grisso, T. (2002). Adolescent development and the measurement of juvenile psychopathy. *Law and Human Behavior*, 26(2), 219-239.
- Skeem, J., & Cauffman, E. (2003). Views of the downward extension: Comparing the youth version of the Psychopathy Checklist with the Youth Psychopathic Traits Inventory. *Behavioral Sciences & the Law*, 21(6), 737-770.
- Snell, T.L., & Morton, D.C. (1991). Survey of state prison inmates, 1991: Women in prison. *Bureau of Justice Statistics Bulletin*.
- Strand, S., & Belfrage, H. (2005). Gender differences in psychopathy in a Swedish offender sample. *Behavioral Sciences & the Law*, 23(6), 837-850.
- Vassileva, J., Kosson, D., Abramowitz, C., & Conrod, P. (2005). Psychopathy versus

- psychopathies in classifying criminal offenders. *Legal and Criminological Psychology*, 10(1), 27-43.
- Verona, E., Hicks, B., & Patrick, C. (2005). Psychopathy and suicidality in female offenders: Mediating influences of personality and abuse. *Journal of Consulting and Clinical Psychology*, 73(6), 1065-1073.
- Vincent, G., & Grisso, T. (2005). A developmental perspective on adolescent personality, psychopathology, and delinquency. *Mental Health Screening and Assessment in Juvenile Justice* (pp. 22-43). New York, NY, US: Guilford Press.
- Vincent, G. M., Vitacco, M. J., Grisso, T., & Corrado, R. R. (2003). Subtypes of adolescent offenders: Affective traits and antisocial behavior patterns. *Behavioral Sciences & the Law*, 21, 695-712.
- Vitale, J. E., & Newman, J. P. (2001). Using the Psychopathy Checklist-Revised with female samples: reliability, validity, and implications for clinical utility. *Clinical Psychology*, 8(1), 117-132.
- Vitale, J., Smith, S., Brinkley, C., & Newman, J. (2002). The reliability and validity of the Psychopathy Checklist--Revised in a sample of female offenders. *Criminal Justice and Behavior*, 29(2), 202-231.
- Warren, J. I., Burnette, M. L., South, S. C., Chauhan, P., Bale, R., Friend, R., Van Patten, I. (2003). Psychopathy in women: Structural modeling and comorbidity. *International Journal of Law and Psychiatry*, 26, 223-242.
- Weiler, B., & Widom, C. (1996). Psychopathy and violent behaviour in abused and neglected young adults. *Criminal Behaviour and Mental Health*, 6(3), 253-271.
- Zaitchik, M.C., & Barese, T.H. (2008). Personality disorders. In B. L. Cutler (Ed.),

Encyclopedia of psychology and law (pp.554-558). Thousand Oaks, CA: Sage Publications.

Appendix A
Psychopathy Checklist: Youth Version
(Forth, Kosson, & Hare, 2003)

Item
1. Impression management
2. Grandiose sense of self-worth
3. Stimulation seeking
4. Pathological lying
5. Manipulation for personal gain
6. Lack of remorse
7. Shallow affect
8. Callous/lack of empathy
9. Parasitic orientation
10. Poor anger control
11. Impersonal sexual behavior
12. Early behavior problems
13. Lacks goals
14. Impulsivity
15. Irresponsibility
16. Failure to accept responsibility
17. Unstable interpersonal relationships
18. Serious criminal behavior
19. Serious violations of conditional release
20. Criminal versatility

Appendix B
Two and Four Factor Structures of PCL:YV
(Forth, Kosson, & Hare, 2003)

Item	2 Factor	4 Factor
1. Impression management	1	1
2. Grandiose sense of self-worth	1	1
3. Stimulation seeking	2	3
4. Pathological lying	1	1
5. Manipulation for personal gain	1	1
6. Lack of remorse	1	2
7. Shallow affect	1	2
8. Callous/lack of empathy	1	2
9. Parasitic orientation	2	3
10. Poor anger control	2	4
11. Impersonal sexual behavior	-	-
12. Early behavior problems	2	4
13. Lacks goals	2	3
14. Impulsivity	2	3
15. Irresponsibility	2	3
16. Failure to accept responsibility	1	2
17. Unstable interpersonal relationships	-	-
18. Serious criminal behavior	2	4
19. Serious violations of conditional release	2	4
20. Criminal versatility	2	4

Factor 1 (interpersonal + affective): Interpersonal/Affective

Factor 2 (behavioral + antisocial): Socially Deviant Lifestyle

F1: Interpersonal: Interpersonal

F2: Affective: Affective

F3: Behavioral: Lifestyle

F4: Antisocial: Antisocial

Appendix C

Subject ID: _____ **Rater:** _____

Item	Score	Factor 1	Factor 2	Factor 3	Factor 4
1. Impression management					
2. Grandiose sense of self-worth					
3. Stimulation seeking					
4. Pathological lying					
5. Manipulation for personal gain					
6. Lack of remorse					

7. Shallow affect					
8. Callous/lacking empathy					
9. Parasitic orientation					
10. Poor anger control					
11. Impersonal sexual behavior					
12. Early behavior problems					
13. Lacks goals					
14. Impulsivity					
15. Irresponsibility					
16. Failure to accept responsibility					
17. Unstable interpersonal relationships					
18. Serious criminal behavior					
19. Serious violations of conditional release					
20. Criminal versatility					
Total Score:					

Omitted Items:					
-----------------------	--	--	--	--	--

Prorated score:					
------------------------	--	--	--	--	--

Total Score:


Factor 1: Interpersonal:

Factor 2: Affective:

Factor 3: Lifestyle:

Factor 4: Antisocial:

Massachusetts Department of Youth Services Institutional Review Board



The Commonwealth of Massachusetts

Executive Office of Health and Human Services

Department of Youth Services

27 Wormwood Street, Suite 400

Boston, MA 02210-1613

617.727.7575

FAX#: 617.951.2409

DEVAL PATRICK
GOVERNOR

TIMOTHY MURRAY
LIEUTENANT GOVERNOR

JUDYANN BIGBY, M.D.
SECRETARY

Frank DiCataldo, Ph.D.
Bedford Policy Institute
35 Braintree Hill Office Park
Braintree, MA 02184

November 20, 2007

Dear Frank:

I am pleased to notify you that your project, *Violence Risk in Adolescent Females* (Principal Investigators: Nathan Cook and Trevor Barese) has been approved by the Institutional Review Board of the Massachusetts Department of Youth Services.

At the completion of your research, please send a copy of the final report to me at the address below. Best of luck with this project. If you have any questions or comments, feel free to contact me.

Sincerely,

Robert Tansi
Institutional Review Board Chair
Department of Youth Services
27 Wormwood St., Suite 400
Boston, MA 02210-1613

Tel: 617-960-3348

Email: robert.tansi@state.ma.us

Appendix E
Forensic Evaluation Data Sheet
(Bedford Policy Institute)

I. Demographic Information

Name:
Age:
DOB:
Date of Commitment:
Mid#:
Area:
Committing Court:
DYS Program:
Dates of Interview:
Name of Evaluator:
Race/Ethnicity:
Gender:

Legal Status: Commit to 18 Youthful Offender Extension of Commit
Detained

Type of Evaluation: Class Extension 68(a) Assess Testing
Number of Commitments:
Referral Number:

II. Delinquency History Information

List of Prior Delinquency Adjudication and Legal Findings:

Name of the Offense Outcome and Date	Date of Arraignment	Legal
-----------------------------------------	---------------------	-------

Commitment offense(s):

Name of the Offense	Date of Arraignment
---------------------	---------------------

III. Mental Health History and Data

Prior psychiatric hospitalization: Yes or No

Number of psychiatric hospitalizations: _____

Current Medication: Yes or No

Name of *current* medications:

Name of *prior* medication:

History of suicide attempts: Yes or No

Number of suicide attempts: _____

Methods Used and #: Overdose (#) Cutting (#) Hanging (#)
Other: _____

History of suicide threats: (only if there is no hx of attempts): Yes or No

Self Injurious Behavior: Yes or No
Scratching Inserting Foreign Objects Ingesting Foreign Objects Head
Banging Burning Other:

Prior Diagnoses:

IV. *Clinical Data/ Risk Factors*

Positive Parental Support or Nurturance: Yes No Not Clear

Parental Control and Accountability for Juvenile: Yes No Not Clear

Hx of attachment problems early childhood: Yes No Not Clear

History of abuse: Yes or No

Type of abuse: Physical Sexual Emotional Neglect

Prior History of DSS Services: Yes or No

Prior History of CHINS: Yes or No

Academic Achievement: High Average Poor No data

History of Truancy: Yes or No

Fighting in School: Yes or No

Disruptive Behavior at School: Yes or No

Weapons at School: Yes or No

Retained a Grade: Yes or No **If yes, how many:** _____

IQ Level: Superior or Above Average Below Average Borderline
MR Unknown

Hx of special education services: Yes or No

Behavior Problems: _____
Learning Disability: _____
Both: _____

Substance abuse problems: Yes or No

Type of Substances Abused:

Negative peer relationships: Yes or No

Gang Affiliation: Yes or No

Pro-social or positive interests or hobbies: Yes or No or Unknown

What are they? _____

Admits to Commitment Offense: Yes Partial No

Blames the Victim: Yes Partial No

Blames external factors: Yes Partial No

Minimizes harm: Yes Partial No

Mode of violence: Reactive Proactive Mixed Unknown N/A

V. Sexual Offense (If commitment offense is not a sexual offense, skip to next section)

Type of victim: Child (5 yrs. Younger) Peer aged Adult Disabled
Mixed

Age of victim: _____

Gender of victim: _____

Relationship to victim: stranger acquaintance girlfriend bio sib
step/foster sib

Location: residence outdoors motor vehicle other: _____

Time: _____

Type of offense: Solitary or Group

Number of co-defendants: _____

History of prior sexual offenses: Yes or No

Number of prior sexual offenses: _____

History of violent delinquency: Yes or No

History of non-violent delinquency: Yes or No

Method of victim compliance: Grooming Threat Force Violence
Other:

Type of sexual assault: Touching Forced oral sex Vaginal Intercourse
Anal intercourse

Weapon present: Yes or No

Type of weapon: _____

Violence Used: Yes or No

Level of victim injury: Mild Moderate Severe

Deviant arousal pattern: Pedophilic Violent other: _____ unknown

Substance abuse at time of offense: Yes or No

► *Violent Offense (if commitment offense is a sexual offense, do not complete this section)*

Type of offense: Solitary or Group

Number of co-defendants: _____

Weapon present: Yes or No

Type of weapon: Handgun Shotgun or rifle Knife Blunt object
other: _____

Victim injury: Yes or No

Level of victim injury: Mild Moderate Severe

Verbal threat: Yes or No

Substance abuse at time of offense: Yes or No

► *Victim Characteristics*

Number of victims: _____

Gender:

Age:

Race:

Relationship: Friend Girl/boyfriend Family member Stranger
Acquaintance Rival

Location: Residence School Outdoors MBTA Public
building

Time: _____

VI. Conclusions

1. Diagnostic Impressions

Diagnoses, including substance abuse:

Recommendation of DMH services: Yes or No

Type of service recommended: Inpatient IRTP Residential
Case management

2. Risk Assessment

Risk factors identified: (Highlight all that apply)

1. Early childhood abuse
2. Witnessed domestic violence
3. Anti-social role modeling
4. Poor attachment history
5. Parental mental illness
6. Parental substance abuse
7. Early developmental/emot. problems
8. Early pattern of undercontrolled behv.
9. Early aggression/destructiveness
10. Poor early peer socialization
11. Poor school functioning
12. Substance abuse
13. Negative peer group
14. Poor parental control
15. Poor parental support/nurturance
16. Weapon possession
17. Violence history
18. Impulsivity/low self-control
19. No pro-social interests
20. Grandiose/self-inflated:
21. Externalizes blame
22. Justifies behavior
23. Minimizes harm
24. Low empathy
25. Thrill seeking
26. Dominance/power needs
27. Depression
28. High harm vigilance
29. Psychotic paranoia
30. Perceives malevolent threat or challenge
31. Violence as means to an end
32. Anger
33. Retaliation
34. Other:_____

Risk level: High Moderate Low

3. *Placement and Treatment Needs*

a. Placement recommendation: Secure Residential Day reporting with
clinical services DMH

b. Treatment needs: (highlight all that apply)

1. Anger control
2. Substance abuse
3. Mental health
4. Sex offender (cog)
5. Sex offender (recondition)
6. Social skill
7. Violence relapse prevention
8. Family therapy
9. Dynamic psychotherapy for trauma/loss
10. Behavioral management
11. Other: _____

I.V. CODING SHEET

Subject ID: _____

DEMOGRAPHICS

Gender of offender *Female* *Male*

Age of offender

Race of offender

VICTIM/OFFENSE

Exact relationship to victim

FED FORM _____

Number of victims

Gender of victim *Female* *Male*

Mode of violence: *Reactive* *Proactive*
Mixed

Relationship: *Friend* *Girl/boyfriend* *Family*
member
Stranger *Acquaintance* *Rival*

Rater: _____

ABUSE/ATTACHMENT

Age at first interruption of family structure

Number of previous living arrangements

FED FORM _____

Positive Parental Support or Nurturance

Yes *No* *Not Clear*

Parental Control and Accountability for Juvenile

Yes *No* *Not Clear*

Hx of attachment problems early childhood

Yes *No* *Not Clear*

History of abuse *Yes* *No*

Type of abuse *Physical* *Sexual*
Emotional *Neglect*

Prior History of DSS Services *Yes* *No*

Witnessed domestic violence *Yes* *No*

Antisocial role modeling *Yes* *No*

Appendix F
I.V. CODING SHEET

FED
FORM

Type of service recommended *Inpatient* *IRTP* *Residential*

Case management

Placement recommendation *Secure Residential* *Day reporting with clinical services*

DMH

Treatment needs:

Anger control

Substance abuse

Mental health

Sex offender (cog)

Sex offender (recondition)

Social skill

Violence relapse prevention

Family therapy

Dynamic psychotherapy for trauma/loss

Behavioral management

Other: _____

Number of prior psychiatric

hospitalizations

Table 1 *Descriptive Statistics for Females (N=50) and Males (N=50)*

	Females			Males		
Race	N	%	Valid N	N	%	Valid N
Caucasian	26	52	50	23	46	50
African American	13	26	50	13	26	50
Hispanic	7	14	50	11	22	50
Asian	2	4	50	1	2	50
Multiracial	2	4	50	2	4	50
Childhood trauma	N	%	Valid N	N	%	Valid N
History of abuse	42	84	50	29	58	50
History of physical abuse	27	54	50	19	38	50
History of neglect	24	48	50	19	38	50
History of emotional abuse	22	44	50	11	22	50
History of sexual abuse	24	48	50	12	24	50
Positive parental support/ nurt.	9	18	38	18	36	39
Parental control or accountability	6	12	45	9	18	42
Witnessed domestic violence	22	44	50	15	30	50
Prior DSS involvement	39	78	50	28	56	50
History of attachment problems	22	44	30	19	38	31
Variables	<i>M</i>	<i>SD</i>	Range	<i>M</i>	<i>SD</i>	Range
Age	15.82	1.19	14 – 19	16.34	1.24	13 - 19
PCL:YV total score	17.97	5.18	6.3 – 30	17.68	6.15	8 – 35
Interpersonal features	3.15	1.97	0 – 8	2.82	2.17	0 – 8
Affective features	4.00	2.03	0 – 8	4.19	2.19	0 – 8
Interpersonal/Affective	7.15	3.22	1 – 15	7.01	3.87	0 – 15.5
Lifestyle features	4.64	1.65	0 – 8	4.43	1.58	1.2 – 8
Antisocial features	4.31	1.81	0 – 7	4.74	1.84	1 – 8.5
Lifestyle/Antisocial	8.95	2.8	2 – 14	9.17	2.69	2.2 – 16.5
Number of previous living arrangements	4.8	5	1-24	4	3.5	0-17
Age at first interruption of family structure	9.3	5.1	5-16	8.9	5.5	0-16

Table 2

Correlation Between Childhood Traumas and PCL:YV Scores

Variable	PCL:YV	
	Low	High
History of Sexual Abuse	28.6%	71.4%**
History of Neglect	34.6%	65.4%*
Prior History of DSS Services	39.0%	61.0%**

Note Low = PCL:YV total < 15 (1/3N), High = PCL:YV total > 21 (1/3N)

* p =.065, **p<.05

Table 3

Correlation Between Parenting Skills and PCL:YV Scores

<i>Variable</i>		<i>PCL:YV</i>	
		Low	High
Female	Positive Parental Support or Nurturance	83.3%*	16.7%
Male	Parental Control and Accountability for Juvenile	100.0%*	.0%

Note Low = PCL:YV total < 15 (1/3N), High = PCL:YV total > 21 (1/3N)

* p<.05

Table 4

Summary of Standard Regression for Variables Predicting PCL:YV Total Score

<i>Variable</i>	<i>B</i>	<i>SE B</i>	<i>β</i>
Parental Control and Accountability for Juvenile	-3.72	1.57	-.25*
Prior History of DSS Services	2.74	1.36	.23*
Number of Previous Living Arrangements	-.16	.15	-.12

Note $R^2 = .12$ ($p < .05$)

* $p < .05$

Table 5

Summary of Standard Regression for Variables Predicting PCL:YV Total Score Between Genders

<i>Gender</i>	<i>Variable</i>	<i>B</i>	<i>SE B</i>	<i>β</i>
Female	Parental Control and Accountability for Juvenile	-3.68	2.32	-.24
	Prior History of DSS Services	1.12	1.94	.09
	Number of Previous Living Arrangements	-.11	.16	-.10
Male	Parental Control and Accountability for Juvenile	-3.42	2.24	-.23
	Prior History of DSS Services	4.68	2.24	.38*
	Number of Previous Living Arrangements	-.32	.32	-.18

Note male $R^2 = .19$ ($p < .05$)

* $p = .043$

Table 6

Summary of Standard Regression for Variables Predicting PCL:YV Factor 1
(Interpersonal/Affective) Score

<i>Gender</i>	<i>Variable</i>	<i>B</i>	<i>SE B</i>	<i>β</i>
Female	Number of Previous Living Arrangements	-.20	.10	-.31*
	Positive Parental Support or Nurturance	-3.42	1.16	-.46**
	History of Sexual Abuse	1.0	.97	.16
Male	Number of Previous Living Arrangements	-.09	.22	-.08
	Positive Parental Support or Nurturance	-.26	1.34	-.03
	History of Sexual Abuse	.35	1.74	.04



Note female $R^2 = .26$ ($p < .05$)






* $p = .056$ ** $p = .006$

Figure Captions

Figure 1. Significant differences between low and high PCL:YV groups in prevalence of distinct forms of childhood abuse.

Extreme High/Low Omit Middle
Low High



-  History of Physical Abuse
-  History of Sexual Abuse
-  History of Emotional Abuse
-  History of Neglect
-  no hx of abuse